

Could light harvesting complexes exhibit non-classical effects at room temperature?

Mark M. Wilde and James M. McCracken

*Electronic Systems Division, Science Applications International Corporation,
4001 North Fairfax Drive, Arlington, Virginia, USA 22203*

Ari Mizel

Laboratory for Physical Sciences, 8050 Greenmead Drive, College Park, Maryland, USA 20740

(Dated: November 5, 2009)

Mounting experimental and theoretical evidence suggests that coherent quantum effects play a role in the efficient transfer of an excitation from a chlorosome antenna to a reaction center in the Fenna-Matthews-Olson protein complex. However, it is conceivable that a satisfying alternate interpretation of the results is possible in terms of a classical theory. To address this possibility, we consider a class of classical theories satisfying the minimal postulates of macrorealism and frame Leggett-Garg-type tests that could rule them out. Our numerical simulations indicate that even in the presence of decoherence, several tests could exhibit the required violations of the Leggett-Garg inequality. Remarkably, some violations persist even at room temperature for our decoherence model.

PACS numbers: 03.65.Ta, 03.65.Yz, 82.39.Jn

Keywords: Leggett-Garg inequality, macrorealism, Fenna-Matthews-Olson protein complex, Prosthecochloris aestuarii, quantum effects in biological systems

I. INTRODUCTION

Does quantum mechanics play a nontrivial role in evolutionary or molecular processes in biological systems? How could this result be true, when biological systems interact in environments that are “hot” and “wet”? Furthermore, how can we frame the first question *rigorously*? These questions have been the subject of intense debate [1, 2] and a surge of recent interest from both the quantum information community [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16] and the experimental quantum chemistry community [17, 18, 19, 20]. This active area is sometimes described as “quantum biology” [2, 11]. Some have suggested that, in addition to the possibility of remarkable new insights into biology, there is the potential to harness naturally occurring coherent quantum effects in biology to engineer new devices.

Two biological processes are playing a particularly important role in fueling interest in this subject: energetic excitation transfer in the Fenna-Matthews-Olson (FMO) light-harvesting protein complex of green bacteria [3, 4, 12, 13, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28] and the radical ion-pair mechanism in the so-called avian compass [6, 8, 9, 10, 14]. For each system, there are claims of quantum mechanical behavior in the scientific literature. On the theory side, several studies in the literature have proposed an open quantum systems approach to model the dynamics of these biological processes [3, 4, 12, 13] and have lent credence to the proposition that coherent quantum effects play a role in their functionality.

Agreement between quantum theoretical models and experiment, however, does not *irrevocably* demonstrate the presence of quantum effects. This point is subtle,

but it is a logical fallacy, called the “affirmation of the consequent,” to conclude once and for all that the quantum biological models are the correct models simply because they coincide with the observations of some experiments (Leggett stresses this point in several of his papers [29, 30]). It would be useful to *irrevocably* exclude certain classes of classical models that might apply to these biological systems by considering fundamental tests of non-classicality.

One might wonder if it is well-motivated to conduct a test for non-classicality on a system such as the FMO complex that is clearly microscopic. In fact, is it not obvious that the FMO complex should behave according to quantum-mechanical laws, given that the length scales and time scales are those to which quantum theory applies? We argue that applying a test for non-classicality is valuable here because a significant amount of environmental noise acts on the chromophores of the FMO complex. These decohering effects may “wash out” quantum-mechanical behavior and make it appear as if the FMO complex behaves according to a class of macrorealistic models (after all, some argue that classical behavior in general arises due to such decoherence [31]). In some cases (in a high temperature limit), researchers actually have modeled the dynamics of the FMO complex with a classical incoherent hopping (Förster) model [21, 22, 32, 33]. Recent results [3, 4, 12, 13] suggest that a classical incoherent hopping model is insufficient to explain the ultra-efficient transfer of energy in photosynthesis, but a test for non-classicality could irrevocably exclude the whole class of models to which the classical incoherent hopping model belongs.

In this paper, we frame a test for non-classicality in the FMO protein complex, the molecular complex responsible for the transfer of energetic excitations in a photosynthetic reaction. Quantum chemists have deter-

mined a tight-binding Hamiltonian for this simple system [24]. Phenomenological modifications to the standard Schrödinger equation have allowed an open quantum systems model of its dynamics [3, 4, 12, 13]. Motivated by experimental results, several theoretical studies have computed the efficiency of energy transfer from the chlorosome antenna to the reaction center in green bacteria and have asserted that coherent quantum effects play a role. To assess this claim, we frame a test for violation of *macrorealism*, as quantified by the Leggett-Garg inequality [34]. In this sense, we are following the program of Leggett outlined in Ref. [29] and the suggestion of Zeilinger in Ref. [1] that it would be useful to subject biomolecules to tests of non-classicality. The Bell [35, 36] and Leggett-Garg [53] tests of non-classicality are benchmarks that determine whether a given dynamical system has stronger-than-classical spatial or temporal correlations, respectively. Each of these tests provides an inequality that bounds the spatial or temporal correlations of a given system—a violation of the inequality implies that the system in question does not behave in a classical manner. The advantage of the Leggett-Garg test over a Bell test is that a Leggett-Garg test applies to a *single* quantum system, easing experimental difficulty. The Leggett-Garg inequality was originally applied to superconducting quantum systems. It was later considered for other systems such as quantum dots [37, 38, 39] and photons [40] but, to our knowledge, this is the first application of the Leggett-Garg theory to a biological system.

We organize the rest of this paper as follows. We introduce the Leggett-Garg inequality in the next section and review a simple example of a two-level quantum system that violates it. In Section III, we review the quantum dynamical model of the FMO complex (we specifically employ the nine-level model of Refs. [3, 4], which is a modification of the seven-level model of Rebentrost *et al.* in Refs. [12, 13, 16]). We then discuss several examples of dichotomic observables for the FMO complex to which we can apply the Leggett-Garg inequality. Section IV begins our study of the Leggett-Garg inequality and the FMO complex. We first study the Leggett-Garg inequality with purely coherent dynamics and are able to derive analytical results. These analytical formulas allow us to determine exactly when coherent dynamics give a violation of the inequality. We then exploit these analytical results in our numerical simulations of the Leggett-Garg inequality and the FMO complex, where we show that it is still possible to violate the inequality even in the presence of noise and potentially even at room temperature. We conclude with some open questions for further study.

II. THE LEGGETT-GARG INEQUALITY

The Leggett-Garg inequality applies to any system that obeys the postulates of macrorealism. The postulates of macrorealism for a two-level system are as follows [29, 34]:

1. *Macrorealism per se*: A macroscopic object is in one of two definite states at any given time.
2. *Noninvasive measurement*: It is possible in principle to determine the state of the system without affecting it or any subsequent dynamics.
3. *Induction*: The properties of ensembles are determined exclusively by initial conditions (and, in particular, not by final conditions).

It is reasonable to assume that a classical system, in principle, should obey the postulates of a macrorealistic theory.

The Leggett-Garg inequality bounds the two-time correlation functions of three dichotomic observables $Q(t_1)$, $Q(t_2)$, and $Q(t_3)$ measured at respective times t_1 , t_2 , and t_3 . The observables $Q(t_1)$, $Q(t_2)$, and $Q(t_3)$ could be the spin of a particle or the location of the trapped magnetic flux in a double-well potential as in the original application of Leggett and Garg [34]. Let $C_{i,j}$ denote the following two-time correlation function:

$$C_{i,j} \equiv \langle Q(t_i) Q(t_j) \rangle.$$

The Leggett-Garg quantity K is the following combination of three two-time correlations and a constant:

$$K \equiv C_{1,2} + C_{2,3} + C_{1,3} + 1. \quad (1)$$

Note that we can obtain the alternate Leggett-Garg quantities

$$-C_{1,2} + C_{2,3} - C_{1,3} + 1, \quad (2)$$

$$-C_{1,2} - C_{2,3} + C_{1,3} + 1, \quad (3)$$

$$C_{1,2} - C_{2,3} - C_{1,3} + 1, \quad (4)$$

merely by flipping the sign of the respective observables $Q(t_1)$, $Q(t_2)$, and $Q(t_3)$. The following Leggett-Garg inequality bounds the Leggett-Garg quantity K when the system in question is a macrorealistic system being measured noninvasively:

$$K \geq 0. \quad (5)$$

The last correlation function $C_{1,3}$ is to be obtained experimentally by measuring at times t_1 and t_3 but refraining from measuring at time t_2 . By comparing $C_{1,3}$ to correlation functions $C_{1,2}$ and $C_{2,3}$ obtained in the presence of a measurement at t_2 , the Leggett-Garg inequality is sensitive to invasiveness in the t_2 measurement.

An example of a system that violates the Leggett-Garg inequality is a spin-1/2 particle with system Hamiltonian $H = \omega \sigma_X / 2$ and with the observables $Q(t_1) = Q(t_2) = Q(t_3) = \sigma_Z$. This choice leads to the following value of the Leggett-Garg quantity K in (1):

$$K = \cos(2\omega\Delta t) + 2\cos(\omega\Delta t) + 1,$$

where the parameter Δt is the uniform time interval between the successive measurements of the observable

Q . Observe that choosing the interval $\Delta t = 3\pi/4\omega$ sets $K = -\sqrt{2}+1$ and leads to a violation of the Leggett-Garg inequality in (5). Thus, this quantum system does not obey the postulates of a macrorealistic theory when we choose the measurement time intervals as given above. This violation is perhaps not surprising because a spin-1/2 system is a genuine quantum system and “cannot have the objective properties tentatively attributed to macroscopic objects prior to and independent of measurements” [41].

III. MODEL FOR THE FMO COMPLEX

Much of the “quantum biological” interest has focused on energy transport in the Fenna-Matthews-Olson (FMO) protein complex [23], which is believed to be the main contributor to ultra-efficient energy transfer in photosynthesis. The FMO protein complex is a trimer in the bacterial species *prothecochloris aestuarii*. The theoretical models in the literature [3, 4, 12, 13, 16] apply to the dynamics of one unit of the trimer. The model assumes that a photon impinges on the peripheral antenna of the light harvesting complex. Absorption of the photon produces an electronic excitation, an exciton, that then traverses a network of seven chromophores or *sites* in one unit of the trimer. The exciton can either recombine, representing a loss of the excitation, or it can transfer to a reaction center, where a light-to-charge conversion occurs before energy storage. Theoretical models [12, 13] indicate that coherent quantum effects combined with decoherence may lead to a quantum stochastic walk [42] that transports energy efficiently. Rebentrost *et al.* provide evidence that coherent quantum effects are responsible for the ultra-high efficiency of photosynthesis [13] by demonstrating that the transport efficiency is much higher with coherent quantum effects than it is without.

Our physical model for excitation transfer is the nine-level model in Refs. [3, 4], a modification of the seven-level model in Refs. [12, 13, 16]. We can restrict dynamics to the single-excitation manifold because the excitation number is a conserved quantity in the absence of light-matter interaction events (within the exciton recombination time scale of 1 ns [44]). The possible states for the exciton can be expressed in the *site basis* $\{|m\rangle\}_{m=1}^7$, where the state $|m\rangle$ indicates that the excitation is present at site m . The incoherent dynamics include a “ground” state $|G\rangle$ corresponding to the loss or recombination of the excitation and a sink state $|S\rangle$ corresponding to the trapping of the exciton at the reaction center. The excitation evolves into one of the two states $|G\rangle$ or $|S\rangle$ in the limit of infinite time. The density operator ρ for this open quantum system admits the following representation in the site basis:

$$\rho = \sum_{m,n \in \{G,1,\dots,7,S\}} \rho_{m,n} |m\rangle \langle n|.$$

We simplify our analysis by assuming that the dy-

namics of the density operator are Markovian. Thus, a Lindblad master equation, with coherent and incoherent components, models the dynamics [45, 46]. Coherent evolution occurs according to the following Hamiltonian H [24]:

$$H \equiv \begin{bmatrix} 215 & -104.1 & 5.1 & -4.3 & 4.7 & -15.1 & -7.8 \\ -104.1 & 220 & 32.6 & 7.1 & 5.4 & 8.3 & 0.8 \\ 5.1 & 32.6 & 0 & -46.8 & 1.0 & -8.1 & 5.1 \\ -4.3 & 7.1 & -46.8 & 125 & -70.7 & -14.7 & -61.5 \\ 4.7 & 5.4 & 1.0 & -70.7 & 450 & 89.7 & -2.5 \\ -15.1 & 8.3 & -8.1 & -14.7 & 89.7 & 330 & 32.7 \\ -7.8 & 0.8 & 5.1 & -61.5 & -2.5 & 32.7 & 280 \end{bmatrix}, \quad (6)$$

where the above matrix representation of H is with respect to the site basis $\{|m\rangle\}_{m=1}^7$, and the units of energy are cm^{-1} (the typical units of choice in spectroscopy experiments). The diagonal terms in H correspond to the site energies, and the off-diagonal terms correspond to intersite couplings. The order of magnitude of the energies in the above Hamiltonian is 100 cm^{-1} , implying that we should observe dynamics on the order of 300 femtoseconds (fs) (Fleming *et al.* experimentally observed behavior on this order in Ref. [17]).

Three Lindblad superoperators [45, 46] also contribute to the dynamics of the density operator in the nine-level model in Refs. [3, 4]. The general form of a Lindblad superoperator $\mathcal{L}(\rho)$ is

$$\mathcal{L}(\rho) \equiv \sum_m \zeta_m (2A_m \rho A_m^\dagger - \{A_m^\dagger A_m, \rho\}),$$

where ζ_m is a rate and A_m is a Lindblad operator [45, 46].

The first Lindblad superoperator $\mathcal{L}_{\text{diss}}$ in our model corresponds to the dissipative recombination of the exciton (loss of energy in the system):

$$\mathcal{L}_{\text{diss}}(\rho) \equiv \sum_{m=1}^7 \Gamma_m (2|G\rangle \langle m| \rho |m\rangle \langle G| - \{|m\rangle \langle m|, \rho\}). \quad (7)$$

An excitation at site $|m\rangle$ recombines with rate Γ_m , and $|G\rangle \langle m|$ is the Lindblad operator that effects this dissipation.

The next Lindblad superoperator $\mathcal{L}_{\text{sink}}(\rho)$ accounts for the trapping of the exciton in the reaction center:

$$\mathcal{L}_{\text{sink}}(\rho) \equiv \Gamma_{\text{sink}} (2|S\rangle \langle 3| \rho |3\rangle \langle S| - \{|3\rangle \langle 3|, \rho\}). \quad (8)$$

The Lindblad superoperator $\mathcal{L}_{\text{sink}}$ includes the operator $|S\rangle \langle 3|$ because evidence suggests that site 3 in the FMO complex plays a crucial role in transferring the exciton to the reaction center [24], where it is later exploited for energy storage.

The final Lindblad superoperator $\mathcal{L}_{\text{deph}}$ accounts for the unavoidable dephasing interaction with the environ-

ment:

$$\mathcal{L}_{\text{deph}}(\rho) \equiv \sum_{m=1}^7 \gamma_m (2|m\rangle\langle m|\rho|m\rangle\langle m| - \{|m\rangle\langle m|, \rho\}), \quad (9)$$

where γ_m is the rate of dephasing at site m . Discussion of this rough treatment of the decoherence appears in Refs. [13, 21, 47].

The following Lindblad quantum master equation governs the evolution of the density operator ρ :

$$\dot{\rho} = -i[H, \rho] + \mathcal{L}_{\text{diss}}(\rho) + \mathcal{L}_{\text{sink}}(\rho) + \mathcal{L}_{\text{deph}}(\rho), \quad (10)$$

where we explicitly see the contribution of the Hamiltonian (6) and the noise superoperators (7-9) to the dynamics, and we implicitly set $\hbar = 1$. Evolution according to the above Lindblad evolution equation is completely positive and trace-preserving (CPTP) for any time [45], and we let $\mathcal{N}_{t,t_0}(\rho)$ denote the induced CPTP map corresponding to the evolution of the density operator ρ from an initial time t_0 to some later time t .

IV. THE LEGGETT-GARG INEQUALITY AND THE FMO COMPLEX

In the forthcoming subsections, we consider the application of the Leggett-Garg inequality to the FMO complex. We first discuss several observables that one might measure in a Leggett-Garg protocol. We then obtain analytical results when the dynamics are purely coherent. These analytical results allow us to determine the time intervals between measurements in a Leggett-Garg protocol that lead to the strongest violation of the inequality. We finally use these time intervals in a numerical simulation of the FMO dynamics that includes the effects of noise. The result is that several observables exhibit a strong violation of the inequality for temperatures below room temperature, and the violation persists in some cases up to room temperature.

A. Observable for the Leggett-Garg Inequality in the FMO Complex

Recall that the Leggett-Garg quantity in (1) involves any three dichotomic observables $Q(t_1)$, $Q(t_2)$, and $Q(t_3)$ measured at respective times t_1 , t_2 , and t_3 .

We have freedom in choosing both the observables that we measure and the times at which we measure them. Perhaps the simplest dichotomic observable that we can construct corresponds to the question [41]:

“Is the system in state $|\psi\rangle$ or not?”

The two-element set of measurement operators for this question are as follows: $\{|\psi\rangle\langle\psi|, I - |\psi\rangle\langle\psi|\}$. We assign the value +1 if the system is in the state $|\psi\rangle$ and the value

−1 otherwise. Let $Q_{|\psi\rangle}$ denote the resulting observable where

$$Q_{|\psi\rangle} \equiv |\psi\rangle\langle\psi| - (I - |\psi\rangle\langle\psi|) = 2|\psi\rangle\langle\psi| - I. \quad (11)$$

We might build a dichotomic observable from states in the *exciton basis*. The exciton basis is the energy eigenbasis $\{|\phi_m\rangle\}_{m=1}^7$ of the Hamiltonian H in (6), where

$$\forall m \quad H|\phi_m\rangle = E_m|\phi_m\rangle.$$

Then the dichotomic observable constructed from an energy eigenstate is $Q_{|\phi_m\rangle}$. Note that observables of this form commute with the Hamiltonian H in (6).

Another possibility is to build the dichotomic observable from the site basis. This type of observable asks the question, *Is the excitation at site m ?* where $m \in \{G, 1, \dots, 7, S\}$. The dichotomic observable constructed from a site state is the *site observable* $Q_{|m\rangle}$. Observables of this form do not commute with the Hamiltonian H in (6).

B. Analytical Results for Coherent Dynamics

Let us first suppose that evolution of the FMO complex is coherent and does not include the noisy Lindblad evolution terms in (7-9). This assumption is unrealistic, but it gives a starting point for understanding the Leggett-Garg inequality and the FMO complex before proceeding with the full-blown evolution in (10).

The general form of the two-time correlation functions appearing in the Leggett-Garg quantity K in (1) are as follows for a coherent evolution:

$$\begin{aligned} C_{1,2} &= \frac{1}{2} \text{Tr} \left[Q(t_2) e^{-iH(t_2-t_1)} \{Q(t_1), \rho_{t_1}\} e^{iH(t_2-t_1)} \right], \\ C_{2,3} &= \frac{1}{2} \text{Tr} \left[Q(t_3) e^{-iH(t_3-t_2)} \{Q(t_2), \rho_{t_2}\} e^{iH(t_3-t_2)} \right], \\ C_{1,3} &= \frac{1}{2} \text{Tr} \left[Q(t_3) e^{-iH(t_3-t_1)} \{Q(t_1), \rho_{t_1}\} e^{iH(t_3-t_1)} \right], \end{aligned}$$

where H is the Hamiltonian in (6), ρ_{t_1} is the initial density operator of the FMO complex, $\rho_{t_2} \equiv e^{-iH(t_2-t_1)}\rho_{t_1}e^{iH(t_2-t_1)}$, and $\{\cdot, \cdot\}$ is the anticommutator. As indicated in Section II, the correlator $C_{1,3}$ characterizes an experiment in which measurements occur at times t_1 and t_3 but no measurement occurs at time t_2 .

Suppose first that we prepare the FMO complex in some state $|\psi\rangle$ and take the Leggett-Garg observable to be $Q_{|\psi\rangle}$ as in (11). Thus, this measurement is a “survival probability” measurement [41, 48, 49]. Suppose further that we measure $Q_{|\psi\rangle}$ at uniform time intervals Δt . A straightforward calculation [48, 49] shows that the Leggett-Garg quantity in (3) is equal to

$$4|\langle\psi|\psi_{2\Delta t}\rangle|^2 - 4\text{Re} \left[(\langle\psi|\psi_{\Delta t}\rangle)^2 \langle\psi_{2\Delta t}|\psi\rangle \right], \quad (12)$$

where $|\psi_t\rangle \equiv e^{-iHt}|\psi\rangle$. Recall that a violation of the Leggett-Garg inequality occurs when the above quantity drops below zero.




FIG. 1: (Color online) The Leggett-Garg quantity as a function of the uniform time interval Δt for all seven sites in the FMO complex when evolution is coherent and the initial state is the classical mixture $\pi_{1,6}$. The units of Δt are picoseconds. A dotted line divides each plot into two regions. Points above a dotted line are in a “no violation” region, while points below are in a “violation” region. The convention is the same in Figures 2, 3, and 4. To show overall behavior, the Δt axis has a coarse scale up to five picoseconds (ps). However, an experimental test would require control of Δt to hundreds of femtoseconds.

ρ_{t_1}	Measurement	Leggett-Garg Quantity
$\pi_{1,6}$	$ 1\rangle$	$2(\langle 1 6_{\Delta t}\rangle ^2 + \langle 1 1_{2\Delta t}\rangle ^2 - \text{Re}\{\langle 1 (1_{\Delta t}\rangle\langle 1_{2\Delta t} + 6_{\Delta t}\rangle\langle 6_{2\Delta t}) 1\rangle\langle 1 1_{\Delta t}\rangle\})$
$\pi_{1,6}$	$ 6\rangle$	$2(\langle 6 1_{\Delta t}\rangle ^2 + \langle 6 6_{2\Delta t}\rangle ^2 - \text{Re}\{\langle 6 (1_{\Delta t}\rangle\langle 1_{2\Delta t} + 6_{\Delta t}\rangle\langle 6_{2\Delta t}) 6\rangle\langle 6 6_{\Delta t}\rangle\})$
$\pi_{1,6}$	$ 2\rangle, \dots, 5\rangle, 7\rangle$	$2(\langle m 1_{\Delta t}\rangle ^2 + \langle m 6_{\Delta t}\rangle ^2 - \text{Re}\{\langle m (1_{\Delta t}\rangle\langle 1_{2\Delta t} + 6_{\Delta t}\rangle\langle 6_{2\Delta t}) m\rangle\langle m m_{\Delta t}\rangle\})$
$ 1\rangle\langle 1 $	$ 1\rangle$	$4 \langle 1 1_{2\Delta t}\rangle ^2 - 4\text{Re}\{\langle 1_{2\Delta t} 1\rangle(\langle 1 1_{\Delta t}\rangle)^2\}$
$ 1\rangle\langle 1 $	$ 2\rangle, \dots, 7\rangle$	$2 \langle m 1_{\Delta t}\rangle ^2 - 4\text{Re}\{\langle m 1_{\Delta t}\rangle\langle 1_{2\Delta t} m\rangle\langle m m_{\Delta t}\rangle\} + 2\text{Re}\{\langle 1_{2\Delta t} m_{\Delta t}\rangle\langle m 1_{\Delta t}\rangle\}$
$ 6\rangle\langle 6 $	$ 6\rangle$	$4 \langle 6 6_{2\Delta t}\rangle ^2 - 4\text{Re}\{\langle 6_{2\Delta t} 6\rangle(\langle 6 6_{\Delta t}\rangle)^2\}$
$ 6\rangle\langle 6 $	$ 1\rangle, \dots, 5\rangle, 7\rangle$	$2 \langle m 6_{\Delta t}\rangle ^2 - 4\text{Re}\{\langle m 6_{\Delta t}\rangle\langle 6_{2\Delta t} m\rangle\langle m m_{\Delta t}\rangle\} + 2\text{Re}\{\langle 6_{2\Delta t} m_{\Delta t}\rangle\langle m 6_{\Delta t}\rangle\}$
π	all sites	$\frac{4}{d} - \frac{8}{d} \langle m m_{\Delta t}\rangle ^2 + \frac{4}{d} \langle m m_{2\Delta t}\rangle ^2$

TABLE I: The first column lists the initial state of the FMO complex. The second column lists the site that we measure in a Leggett-Garg protocol. The third column gives the analytical form of the corresponding Leggett-Garg quantity as a function of the uniform time interval Δt when dynamics are purely coherent. We use these formulas to compute the results of Figure 1 and Table II.

If the state $|\psi\rangle$ is an eigenstate $|\phi_m\rangle$ in the exciton basis we should not expect to violate the Leggett-Garg inequality because the observable $Q_{|\phi_m\rangle}$ commutes with the Hamiltonian. We can confirm this intuition by plugging the eigenstate $|\phi_m\rangle$ into (12). Doing so gives a value of zero for (12), thereby saturating the Leggett-Garg inequality, but yielding no violation.

Thus, we set the Leggett-Garg observable to a site observable $Q_{|m\rangle}$ where $m \in \{1, \dots, 7\}$. Sites 1 and 6 are

the chromophores that are closest to the chlorosome antenna and are thus most likely to be the initial state of the FMO complex [13]. The initial state ρ_{t_1} of the FMO complex can be a pure state $|1\rangle\langle 1|$ or $|6\rangle\langle 6|$ or a uniform classical mixture of sites $|1\rangle\langle 1|$ and $|6\rangle\langle 6|$:

$$\pi_{1,6} \equiv \frac{1}{2}(|1\rangle\langle 1| + |6\rangle\langle 6|). \quad (13)$$

One might also consider setting the initial state to the

Initial State ρ_0	Site	K	Δt (ps)
$\pi_{1,6}$	1	-0.25053	0.16678
$\pi_{1,6}$	2	-0.22321	0.16678
$\pi_{1,6}$	3	-0.016389	3.1021
$\pi_{1,6}$	4	-0.01574	1.1008
$\pi_{1,6}$	5	-0.12782	0.13343
$\pi_{1,6}$	6	-0.17994	0.16678
$\pi_{1,6}$	7	-0.094719	0.70048
$ 1\rangle\langle 1 $	1	-0.4935	0.16678
$ 1\rangle\langle 1 $	2	-0.44335	0.16678
$ 1\rangle\langle 1 $	3	-0.065461	3.1355
$ 1\rangle\langle 1 $	4	-0.091838	1.7345
$ 1\rangle\langle 1 $	5	-0.08013	2.2015
$ 1\rangle\langle 1 $	6	-0.0097707	0.16678
$ 1\rangle\langle 1 $	7	-0.085607	1.034
$ 6\rangle\langle 6 $	1	-0.0077476	0.13343
$ 6\rangle\langle 6 $	2	-0.0043891	1.034
$ 6\rangle\langle 6 $	3	-0.0032073	0.13343
$ 6\rangle\langle 6 $	4	-0.034082	1.4677
$ 6\rangle\langle 6 $	5	-0.27786	4.9701
$ 6\rangle\langle 6 $	6	-0.35011	0.16678
$ 6\rangle\langle 6 $	7	-0.18045	0.70048
π	all	0	all times

TABLE II: The first column lists the initial state of the FMO complex. The second column lists the site observable that the Leggett-Garg protocol measures. The third column lists the strongest violations for each site observable, and the fourth column gives the corresponding time interval Δt that leads to this violation. (For comparison, the strongest possible violation of the inequality is -0.5 .) We obtained these values assuming that evolution is coherent (though, we examined times up to $\Delta t = 5$ ps only). The last row in the table indicates that we do not obtain a violation for any site observable when the initial state is the maximally mixed state.

maximally mixed state

$$\pi \equiv \frac{1}{7} (|1\rangle\langle 1| + \dots + |7\rangle\langle 7|).$$

Table I lists exact expressions for the Leggett-Garg quantity in (3) for these different cases.

Figure 1 plots the Leggett-Garg quantity for each site observable as a function of the uniform time interval Δt , when the initial state is $\pi_{1,6}$. The result is that each of the seven site measurements gives a violation of the Leggett-Garg inequality for some intervals Δt when the dynamics are purely coherent. These results may not be particularly surprising [48, 49] given that the system is quantum, the measurements are sharp, and the dynamics are purely coherent.

Table II lists the time intervals that lead to the strongest violation for $\Delta t \in [0, 5]$ ps. We only consider times up to five picoseconds (ps) because it is likely that

the exciton will trap by this time for the case of incoherent dynamics in the next section. Table II also lists the time intervals that lead to the strongest violation when the initial state is $|1\rangle\langle 1|$, $|6\rangle\langle 6|$, or π . We now use these time intervals for the more realistic case, where the excitation in the FMO complex experiences the noisy contributions in (7-9) from dissipation and dephasing.

C. Numerical Results for Incoherent Dynamics

The Lindblad evolution in (10) is a more realistic model for evolution of the excitation in the FMO complex. It incorporates the effects of excitonic recombination, trapping to the reaction center, and environmental dephasing noise, albeit in a Markovian context. The two-time correlation functions appearing in the Leggett-Garg quantity K in (1) must now be evaluated using the following more general forms:

$$\begin{aligned} C_{1,2} &= \frac{1}{2} \text{Tr} [Q(t_2) \mathcal{N}_{t_2,t_1} (\{Q(t_1), \rho\})], \\ C_{2,3} &= \frac{1}{2} \text{Tr} [Q(t_3) \mathcal{N}_{t_3,t_2} (\{Q(t_2), \mathcal{N}_{t_2,t_1}(\rho)\})], \\ C_{1,3} &= \frac{1}{2} \text{Tr} [Q(t_3) \mathcal{N}_{t_3,t_1} (\{Q(t_1), \rho\})], \end{aligned}$$

where $\mathcal{N}$ is the superoperator that propagates the density operator forward in time according to the evolution in (10).

In order to perform a numerical simulation for the incoherent dynamics (10), we need to fix the trapping rate Γ_{sink} , the recombination rates Γ_m , and the dephasing rates γ_m . We take these rates from Refs. [3, 4]. Unless otherwise stated, the trapping rate $\Gamma_{\text{sink}} = 62.8/1.88 \text{ cm}^{-1}$, corresponding to about $2\pi \cdot c \cdot \Gamma_{\text{sink}} \approx 6 \text{ ps}^{-1}$, where c is the speed of light in units of $\text{cm} \cdot \text{ps}^{-1}$. The recombination rates Γ_m are uniform for all sites in the FMO complex so that $\Gamma_m = \Gamma_{\text{recomb}} = 1/(2 \cdot 188) \text{ cm}^{-1}$, corresponding to about $2\pi \cdot c \cdot \Gamma_{\text{recomb}} \approx 5 \times 10^{-4} \text{ ps}^{-1}$. We assume that the dephasing rate is uniform so that $\gamma_m = \gamma$ for all sites m .

As in the previous subsection, we choose the Leggett-Garg observable to be a site observable $Q_{|m\rangle}$ where $m \in \{1, \dots, 7\}$. The initial state ρ_{t_1} can either be the pure state $|1\rangle\langle 1|$, $|6\rangle\langle 6|$, or a uniform classical mixture of sites $|1\rangle$ and $|6\rangle$ as in (13), for the same reasons mentioned in the previous subsection (we do not consider the maximally mixed state because there is no phenomenological evidence for this case, and furthermore, it does not give a violation even in the coherent case). The time interval Δt between measurements of the Leggett-Garg observable is taken from Table II. Figures 2, 3, and 4 display the Leggett-Garg quantity as a function of the dephasing parameter γ for each site observable $Q_{|m\rangle}$, where $m \in \{1, \dots, 7\}$.

The figures demonstrate that several site observables exhibit a violation even as γ increases. The temperature analysis of the environment in Ref. [13] indicates that a




FIG. 2: (Color online) The above figure displays the Leggett-Garg quantity as a function of the dephasing parameter γ for each site observable $Q_{|m\rangle}$, where $m \in \{1, \dots, 7\}$. The site observables $Q_{|3\rangle}$ and $Q_{|4\rangle}$ do not give a violation of the Leggett-Garg inequality for any amount of dephasing noise, even though we observed a violation in Figure 1 with purely coherent dynamics. Site observables $Q_{|1\rangle}$, $Q_{|2\rangle}$, $Q_{|5\rangle}$, $Q_{|6\rangle}$, and $Q_{|7\rangle}$ give a strong violation of the inequality for low dephasing noise. Each of their corresponding Leggett-Garg quantities has a smooth, monotonic transition to the “no violation” region for stronger dephasing noise, with the Leggett-Garg quantity for site observables $Q_{|1\rangle}$, $Q_{|2\rangle}$, $Q_{|5\rangle}$, $Q_{|6\rangle}$ withstanding the strongest amount of dephasing noise before they make a transition to the “no violation” region. Using the temperature analysis of the environment in Ref. [13], a dephasing rate of 2.1 ps^{-1} corresponds to a temperature of around 77°K and a dephasing rate of 9.1 ps^{-1} corresponds to a temperature of 298°K (room temperature). Site observables $Q_{|1\rangle}$, $Q_{|2\rangle}$, $Q_{|5\rangle}$, and $Q_{|6\rangle}$ give a violation for room temperature, with the Leggett-Garg quantity respectively equal to -0.0039 , -0.0015 , -0.0059 , and -0.0079 . It might be difficult for an experimentalist to observe these violations at room temperature given that they are weak.

dephasing rate of 2.1 ps^{-1} corresponds to a temperature of around 77°K and a dephasing rate of 9.1 ps^{-1} corresponds to a temperature of 298°K (room temperature). Under the assumption that this is approximately correct, Figures 2, 3, and 4 predict a violation at both temperatures. However, it might be difficult for an experimentalist to observe these violations at room temperature given that they are weak.

We have verified the robustness of these predictions against variations in the dynamical parameters in (10). Ref. [24] mentions that the site energies of their calculated Hamiltonian are accurate within $\pm 2 \text{ cm}^{-1}$. We therefore conducted several simulations that added independent, zero-mean Gaussian noise with variance 2 to each entry in the Hamiltonian matrix in (6) to deter-




FIG. 3: (Color online) This figure is similar to Figure 2, with the exception that the initial state of the FMO complex is a pure state at site 1. Site observables $Q_{|1\rangle}$ and $Q_{|2\rangle}$ give a violation for room temperature, with the Leggett-Garg quantity respectively equal to -0.0077 and -0.003 .




FIG. 4: (Color online) This figure is similar to Figure 2, with the exception that the initial state of the FMO complex is a pure state at site 6. Site observable $Q_{|6\rangle}$ gives a violation for room temperature, with the Leggett-Garg quantity equal to -0.0155 .

mine if the violations would still hold under this slight perturbation. The result is that all room temperature violations in Figures 2, 3, and 4 still hold, and in fact, the values of the Leggett-Garg quantities are the same up to the fourth decimal place. The trapping rates in the literature vary substantially, including 0.25 ps^{-1} [24], 1 ps^{-1} [12], or 4 ps^{-1} [16], so we have checked our results for all of these choices. The plots for all these cases are similar

to Figures 2, 3, and 4 and have approximately the same values for violations at room temperature. This finding is reasonable given that most of our measurement times are less than the average trapping times corresponding to these other trapping rates.

Our numerical simulations demonstrate that several choices of measurements lead to a violation of the Leggett-Garg inequality even when noise processes act on an excitation in the FMO complex. Experimental confirmation could irrevocably exclude a class of macrorealistic theories from describing the dynamics of the excitation.

V. CONCLUSIONS

We have framed tests that could be used to experimentally exclude a class of macrorealistic theories, including a classical incoherent hopping model, from describing the room-temperature dynamical behavior of an excitation in the FMO complex. To do so, we have introduced several examples of observables that one might apply in a test of macrorealism, and our numerical simulations predict that these observables lead to a violation of the Leggett-Garg inequality.

We discuss several ideas for furthering this research. The Leggett-Garg measurements that we have considered here are projective measurements. In practice, an experimentalist is unlikely to realize such an idealized measurement. An experimentalist is more likely able only to effect a “noisy” or “fuzzy” measurement of the sites in the FMO complex. Along these lines, Kofler *et al.* have shown that “fuzzy” measurements may lead to our observation of a classical world even in the presence of quantum dynamics[48, 49]. Future work should

formulate Leggett-Garg inequalities involving quantities that are conveniently accessible to FMO experimentalists (measurements in the site basis are currently infeasible [50]) and should consider the limits of experimental measurement capabilities. Furthermore, future work should consider more realistic models of noise in the FMO complex, potentially including correlated noise [25, 51] and non-Markovian effects [26].

One might consider an application of Bell’s inequality to study the ability of the FMO complex to preserve entanglement.

We have applied the Leggett-Garg inequality to the FMO complex, and it would be valuable to apply the Leggett-Garg test to the dynamics of magnetoreception in the avian compass [6, 8, 9, 10]. It would also be interesting to explore the Leggett-Garg inequality in artificial quantum networks with particular noisy interactions. A study of this sort might lead to an increased understanding of the dividing line between macrorealism and non-classicality in more complicated systems. One might be able to apply the ideas in Refs. [3, 4, 48, 49] here.

Acknowledgments

The authors thank Andrew Cross, Alan Aspuru-Guzik, Kevin Obenland, Patrick Rebentrost, and Leslie Vogt for useful discussions and thank Science Applications International Corporation for supporting this scientific research. MMW is especially grateful to Alan Aspuru-Guzik and Patrick Rebentrost for their warm hospitality during his visit to Harvard and Leslie Vogt for her suggestion to consider Leggett-Garg measurement time intervals within the average trapping time of an exciton.

-
- [1] Derek Abbott, Julio Gea-Banacloche, Paul C. W. Davies, Stuart Hameroff, Anton Zeilinger, Jens Eisert, Howard Wiseman, and Sergey M. Bezrukov. Plenary debate: Quantum effects in biology: Trivial or not? *Fluctuation and Noise Letters*, 8(1):C5–C26, 2008.
 - [2] Derek Abbott, Paul C. W. Davies, and Arun K. Pati, editors. *Quantum Aspects of Life*. Imperial College Press, 2008.
 - [3] Filippo Caruso, Alex W. Chin, Animesh Datta, Susana F. Huelga, and Martin B. Plenio. Highly efficient energy excitation transfer in light-harvesting complexes: The fundamental role of noise-assisted transport. *arXiv:0901.4454*, 2009.
 - [4] Martin B. Plenio and Susana F. Huelga. Dephasing-assisted transport: quantum networks and biomolecules. *New Journal of Physics*, 10(11):113019, 2008.
 - [5] Hans J. Briegel and Sandu Popescu. Entanglement and intra-molecular cooling in biological systems? — a quantum thermodynamic perspective. *arXiv:0806.4552*, 2008.
 - [6] Jianming Cai, Gian Giacomo Guerreschi, and Hans J. Briegel. Quantum control and entanglement in a chemical compass. *arXiv:0906.2383*, 2009.
 - [7] Jianming Cai, Sandu Popescu, and Hans J. Briegel. Dynamic entanglement in oscillating molecules. *arXiv:0809.4906*, 2008.
 - [8] Iannis K. Kominis. Quantum Zeno effect explains magnetic-sensitive radical-ion-pair reactions. *arXiv:0806.0739*, 2008.
 - [9] Iannis K. Kominis. Quantum theory of radical-ion-pair recombination: A new physical paradigm for low-magnetic-field effects. *arXiv:0805.3081*, 2008.
 - [10] Iannis K. Kominis. Quantum Zeno effect underpinning the radical-ion-pair mechanism of avian magnetoreception. *arXiv:0804.2646*, 2008.
 - [11] Seth Lloyd. A quantum of natural selection. *Nature Physics*, 5(3):164–166, March 2009.
 - [12] Masoud Mohseni, Patrick Rebentrost, Seth Lloyd, and Alán Aspuru-Guzik. Environment-assisted quantum walks in photosynthetic energy transfer. *The Journal of Chemical Physics*, 129(17):174106, 2008.
 - [13] Patrick Rebentrost, Masoud Mohseni, Ivan Kassal, Seth Lloyd, and Alán Aspuru-Guzik. Environment-assisted quantum transport. *New Journal of Physics*, 11(3):033003 (12pp), 2009.

- [14] Elisabeth Rieper, Erik Gauger, John J. L. Morton, Simon C. Benjamin, and Vlatko Vedral. Quantum coherence and entanglement in the avian compass. *arXiv:0906.3725*, 2009.
- [15] Joel Gilmore and Ross H. McKenzie. Quantum dynamics of electronic excitations in biomolecular chromophores: Role of the protein environment and solvent. *The Journal of Physical Chemistry A*, 112(11):2162–2176, March 2008.
- [16] Alexandra Olaya-Castro, Chiu Fan Lee, Francesca Fassioli Olsen, and Neil F. Johnson. Efficiency of energy transfer in a light-harvesting system under quantum coherence. *Physical Review B*, 78(8):085115, 2008.
- [17] Gregory S. Engel, Tessa R. Calhoun, Elizabeth L. Read, Tae-Kyu Ahn, Tomá Manal, Yuan-Chung Cheng, Robert E. Blankenship, and Graham R. Fleming. Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature*, 446:782–786, 2007.
- [18] Hohjai Lee, Yuan-Chung Cheng, and Graham R. Fleming. Coherence dynamics in photosynthesis: Protein protection of excitonic coherence. *Science*, 316:1462, 2007.
- [19] Thorsten Ritz, Peter Thalau, John B. Phillips, Roswitha Wiltshcko, and Wolfgang Wiltshcko. Resonance effects indicate a radical-pair mechanism for avian magnetic compass. *Nature*, 429:177–180, 2004.
- [20] Yuan-Chung Cheng and Graham R. Fleming. Dynamics of light harvesting in photosynthesis. *Annual Review of Physical Chemistry*, 60(1):241–262, 2009. PMID: 18999996.
- [21] Jan A. Leegwater. Coherent versus incoherent energy transfer and trapping in photosynthetic antenna complexes. *Journal of Physical Chemistry*, 100:14403–14409, 1996.
- [22] Mirianas Chachisvilis, Oliver Kühn, Tõnu Pullerits, and Villy Sundström. Excitons in photosynthetic purple bacteria: Wavelike motion or incoherent hopping? *Journal of Physical Chemistry B*, 101(37):7275–7283, 1997.
- [23] R. E. Fenna and B. W. Matthews. Chlorophyll arrangement in a bacteriochlorophyll protein from chlorobium limicola. *Nature*, 258(5536):573–577, December 1975.
- [24] Julia Adolphs and Thomas Renger. How proteins trigger excitation energy transfer in the FMO complex of green sulfur bacteria. *Biophysical Journal*, 91(8):2778–2797, 2006.
- [25] Francesca Fassioli, Ahsan Nazir, and Alexandra Olaya-Castro. Multichromophoric energy transfer under the influence of a correlated environment. *arXiv:0907.5183*, 2009.
- [26] Patrick Rebentrost, Rupak Chakraborty, and Alan Aspuru-Guzik. Non-Markovian quantum jumps in excitonic energy transfer. *arXiv:0908.1961*, 2009.
- [27] Herbert Van Amerongen, Leonas Valkunas, and Rienk Van Grondelle. *Photosynthetic Excitons*. World Scientific, 2000.
- [28] Thomas Renger, Volkhard May, and Oliver Khn. Ultrafast excitation energy transfer dynamics in photosynthetic pigment-protein complexes. *Physics Reports*, 343(3):137 – 254, 2001.
- [29] Anthony J. Leggett. Testing the limits of quantum mechanics: motivation, state of play, prospects. *Journal of Physics: Condensed Matter*, 14(15):R415–R451, 2002.
- [30] Anthony J. Leggett. Realism and the physical world. *Reports on Progress in Physics*, 71(2):022001, 2008.
- [31] Wojciech H. Zurek. Decoherence and the transition from quantum to classical. *Physics Today*, 44:36–44, 1991.
- [32] Th. Förster. Energiewanderung und fluoreszenz. *Naturwissenschaften*, 33(6):166–175, June 1946.
- [33] V. M. Kenkre and P. Reineker. *Exciton dynamics in molecular crystals and aggregates*. Springer-Verlag, New York, 1982.
- [34] Anthony J. Leggett and Anupam Garg. Quantum mechanics versus macroscopic realism: Is the flux there when nobody looks? *Physical Review Letters*, 54:857, 1985.
- [35] John S. Bell. On the Einstein Podolsky Rosen paradox. *Physics*, 1:195–200, 1964.
- [36] John S. Bell. *Speakable and Unspeakable in Quantum Mechanics*. Cambridge University Press, 1987.
- [37] Andrew N. Jordan, Alexander N. Korotkov, and Markus Büttiker. Leggett-Garg inequality with a kicked quantum pump. *Physical Review Letters*, 97(2):026805, 2006.
- [38] Nathan S. Williams and Andrew N. Jordan. Weak values and the Leggett-Garg inequality in solid-state qubits. *Physical Review Letters*, 100(2):026804, 2008.
- [39] R. Ruskov, A. N. Korotkov, and A. Mizel. Signatures of quantum behavior in single-qubit weak measurements. *Physical Review Letters*, 96:200404, 2006.
- [40] M. E. Goggin, M. P. Almeida, M. Barbieri, B. P. Lanyon, J. L. O’Brien, A. G. White, and G. J. Pryde. Violation of the Leggett-Garg inequality with weak measurements of photons. *arXiv:0907.1679*, 2009.
- [41] Asher Peres. *Quantum Theory: Concepts and Methods*. Springer, 1st edition, September 1995.
- [42] César A. Rodríguez-Rosario, James D. Whitfield, and Alán Aspuru-Guzik. Quantum stochastic walks: A generalization of classical random walks and quantum walks. *arXiv:0905.2942*, 2009.
- [43] Viv Kendon. Decoherence in quantum walks — a review. *Mathematical Structures in Computer Science*, 17(06):1169–1220, 2007.
- [44] T. G. Owens, S. P. Webb, L. Mets, R. S. Alberte, and G. R. Fleming. Antenna size dependence of fluorescence decay in the core antenna of photosystem I: estimates of charge separation and energy transfer rates. *Proceedings of the National Academy of Sciences of the United States of America*, 84(6):1532–1536, 1987.
- [45] Heinz-Peter Breuer and Francesco Petruccione. *The Theory of Open Quantum Systems*. Oxford University Press, 2007.
- [46] G. Lindblad. On the generators of quantum dynamical semigroups. *Communications in Mathematical Physics*, 48(2):119–130, June 1976.
- [47] H. Haken and G. Strobl. An exactly solvable model for coherent and incoherent exciton motion. *Zeitschrift für Physik A*, 262(2):135–148, April 1973.
- [48] Johannes Kofler and Časlav Brukner. Conditions for quantum violation of macroscopic realism. *Physical Review Letters*, 101(9):090403, 2008.
- [49] Johannes Kofler. *Quantum violation of macroscopic realism and the transition to classical physics*. PhD thesis, University of Vienna, Vienna, Austria, June 2008.
- [50] Alán Aspuru-Guzik and Patrick Rebentrost. private communication, 2009.
- [51] Ahsan Nazir. Correlation-dependent coherent to incoherent transitions in resonant energy transfer dynamics. *arXiv:0906.0592*, 2009.
- [52] D. Home and S. Sengupta. Bell’s inequality and non-

contextual dispersion-free states. *Physics Letters A*,
102(4):159 – 162, 1984.

[53] Leggett points out [29] that single-system Bell-type in-

equalities [34] appeared earlier in Ref. [52].